WORLD JOURNAL OF PHARMACY AND BIOTECHNOLOGY
E-ISSN: 2349-9087 | https://doi.org/10.30904/j.wjpbt.2026.4958

Gastrointestinal Motility Patterns: Mechanisms, Disorders, and Emerging Therapeutic Targets

Dr. SK. Salma Sultana[1]*, Dr. SK. Salma[2], P. Likitha[3], D. Venkata Harsha Vardan[3], Damai Ashok[3]

[1]Professor, Department of Pharmacology, Narayana Pharmacy College, Nellore
[2]Professor, Department of Pharmaceutical Chemistry, Narayana Pharmacy College, Nellore
[3]B Pharmacy student, Narayana Pharmacy College, Nellore
***Corresponding E-mail:** salmasulthanacology@gmail.com*

KEYWORDS
Gastrointestinal motility;
Enteric nervous system;
Peristalsis;
Gut-brain axis;
Motility disorders;
Prokinetics;

Received 30-01-2026
Revised 24-02-2026
Accepted 28-03-2026
Published 21-04-2026

ABSTRACT
Gastrointestinal (GI) motility is a complex, coordinated process essential for digestion, nutrient absorption, immune regulation, and waste elimination. Proper motility depends on the integrated function of smooth muscle layers, the enteric nervous system (ENS), hormonal signaling, and interstitial cells of Cajal (ICC). Disruption of these mechanisms results in clinically significant motility disorders such as achalasia, gastroparesis, irritable bowel syndrome (IBS), and chronic intestinal pseudo-obstruction. Advances in diagnostic tools including high-resolution manometry, wireless motility capsules, electrogastrography, and imaging modalities have enhanced understanding of physiological and pathological motility patterns. Emerging research highlights the importance of the gut-brain axis, immune signaling, microbiome interactions, and molecular pathways in regulating gastrointestinal function. This review summarizes normal motility patterns, regulatory mechanisms, pathophysiology of motility disorders, current pharmacological approaches, and future therapeutic targets.

INTRODUCTION

The gastrointestinal tract performs essential physiological functions including digestion, nutrient absorption, water-electrolyte balance, immune defense, and hormone production. Efficient intestinal motility is critical for these processes. Coordinated movement ensures mechanical breakdown of food, mixing with digestive secretions, controlled transit, and waste elimination. The enteric nervous system (ENS), often referred to as the "second brain," contains more neurons than the spinal cord, autonomously regulates motility patterns. Dysregulation of neural, muscular, or hormonal mechanisms can lead to motility disturbances. Motility disorders are common yet under-recognized in clinical practice. Critically ill patients, postoperative individuals, and patients with neurological diseases frequently develop impaired gastrointestinal motility. Delayed gastric emptying occurs in nearly 50% of mechanically ventilated patients and up to 80% of patients with head injury. Such disturbances increase the risk of bacterial translocation, sepsis, and multiple organ failure. Understanding the mechanisms underlying gastrointestinal motility is essential for improving diagnosis and therapeutic strategies.

2. ANATOMY AND PHYSIOLOGY OF THE GASTROINTESTINAL TRACT

The GI tract wall consists of four main layers:

Mucosa: Responsible for absorption and sensory detection

Submucosa: Contains blood vessels, lymphatics, submucosal (Meissner's) plexus

Muscularis externa: Inner circular and outer longitudinal smooth muscle layers responsible for motility

Serosa: Protective outer covering

The ENS, comprising the myenteric (Auerbach's) and submucosal plexuses, coordinates muscle contraction, secretion, and blood flow.

3. ROLE OF SMOOTH MUSCLE AND NEURAL CONTROL

3.1 Smooth Muscle

Gastrointestinal smooth muscle contracts rhythmically to produce propulsion and mixing:

- **Peristalsis** – Propulsive contractions
- **Segmentation** – Mixing contractions
- **Tonic contractions** – Sustained contractions (e.g., sphincters)

Smooth muscle activity is involuntary and modulated by neural and hormonal inputs.

3.2 NEURAL REGULATION

The autonomic nervous system regulates motility through:

- **Parasympathetic system** – Stimulates motility (via vagus nerve)
- **Sympathetic system** – Inhibits motility and tightens sphincters
- **Enteric nervous system (ENS)** – Coordinates intrinsic motility independently

Key neurotransmitters include:

- Acetylcholine (excitatory)
- Serotonin (5-HT)
- Nitric oxide (inhibitory)
- Dopamine

4. GASTROINTESTINAL MOTILITY PATTERNS

4.1 Peristalsis: Wave-like contractions that propel luminal contents forward.
4.2 Segmentation: Localized contractions that mix chyme with digestive enzymes.
4.3 Migrating Motor Complex (MMC): Cyclic fasting pattern that clears residual contents between meals.
4.4 Tonic Contractions
Sustained contractions in sphincters controlling directional flow.
4.5 Mass Movements
Strong colonic contractions that move fecal material toward the rectum.
4.6 Retropulsion: Backward propulsion in the stomach aiding mechanical digestion.

5. HORMONAL REGULATION OF MOTILITY

Key hormones include:

- **Gastrin** – Enhances gastric motility
- **Cholecystokinin (CCK)** – Slows gastric emptying
- **Secretin** – Reduces gastric motility
- **Motilin** – Initiates MMC during fasting
- **Somatostatin** – Inhibits motility
- **Serotonin (5-HT)** – Promotes peristalsis

6. METHODS OF STUDYING GASTROINTESTINAL MOTILITY

6.1 Imaging Techniques

- Barium studies
- Scintigraphy
- MRI
- Ultrasound

6.2 Electrophysiological Techniques

- Electrogastrography (EGG)
- Intracellular electrode recordings

6.3 Advanced Diagnostic Tools

- Wireless motility capsule
- High-resolution manometry
- Impedance planimetry (EndoFLIP)
- 3D motility mapping

7. PHYSIOLOGICAL VARIATIONS IN MOTILITY

7.1 Segmental Differences

- Esophagus – Rapid peristalsis
- Stomach – Grinding and controlled emptying
- Small intestine – Segmentation + MMC
- Colon – Water absorption + mass movements

7.2 Fasting vs. Fed State

- Fasting – MMC activity
- Fed – Mixing and absorption-focused motility

7.3 Age-Related Changes

- Reduced sphincter tone
- Slower gastric emptying
- Increased constipation prevalence

8. DISORDERS OF GASTROINTESTINAL MOTILITY

Common conditions include:

- Achalasia
- Gastroesophageal reflux disease (GERD)
- Gastroparesis
- Irritable bowel syndrome (IBS)
- Chronic intestinal pseudo-obstruction

8.1 Pathophysiology

Motility disorders result from:

- ENS dysfunction
- Smooth muscle abnormalities
- Sphincter dysfunction
- Hormonal imbalance
- Inflammatory mediators
- Genetic mutations
- Interstitial cells of Cajal dysfunction

9. PHARMACOLOGICAL MODULATION OF MOTILITY

9.1 Prokinetic Agents

- Metoclopramide
- Domperidone
- Erythromycin (motilin agonist)

9.2 Laxatives

Increase colonic transit.

9.3 Antidiarrheal Agents

- Loperamide (reduces motility)

9.4 Antispasmodics

- Hyoscine
- Dicyclomine

9.5 Other Drugs Affecting Motility

- Opiates (reduce motility)
- Antidepressants
- Calcium channel blockers
- Anticholinergic drugs

10. EMERGING RESEARCH AND FUTURE DIRECTIONS

10.1 Novel Diagnostic Advances

- High-resolution manometry
- Wireless motility capsules
- Functional MRI
- Biomarker development

10.2 Molecular and Neural Advances

- Interstitial cells of Cajal research
- Gut-brain axis studies
- Cytokine-mediated modulation
- Microbiome-motility interactions

10.3 Potential Therapeutic Targets

- Serotonin receptor modulators
- Motilin and ghrelin agonists
- Ion channel modulators
- Stem cell therapies
- Microbiota-directed treatments
- Vagus nerve stimulation

11. CLINICAL SIGNIFICANCE

Understanding gastrointestinal motility patterns is crucial for diagnosing and managing functional and structural disorders. Accurate identification of motility disturbances guides therapeutic interventions including prokinetic therapy, nutritional strategies, neuromodulation, and surgical correction.

12. AREAS REQUIRING FURTHER RESEARCH

Future investigations should focus on:

- Molecular mechanisms of ENS regulation
- Personalized medicine approaches
- Microbiome-based therapies
- Safer prokinetic drugs
- Non-invasive diagnostic technologies
- Gene and regenerative therapies

RESULTS AND DISCUSSION

Normal motility includes peristalsis, segmentation, migrating motor complexes, and mass movements. Regulation occurs via the enteric nervous system, autonomic inputs, hormones, and pacemaker activity of interstitial cells of Cajal. Disorders arise from neural dysfunction, smooth muscle impairment, inflammatory mediators, or receptor abnormalities.

Figure 1: Layers of the Gastrointestinal Wall



Figure 2: Peristaltic Movement



Figure 3: Neural and Hormonal Regulation



CONCLUSION

Gastrointestinal motility is a highly regulated physiological process involving coordinated neural, muscular, hormonal, and immune mechanisms. Disruption of this balance results in significant morbidity. Advances in neurogastroenterology, molecular biology, and biomedical technology are transforming our understanding of motility disorders, enabling development of targeted therapies. Continued translational and clinical research is essential to improve patient outcomes.

ACKNOWLEDGMENT

The authors thank the Narayana Pharmacy College, Nellore, Andhra Pradesh, India, for technical assistance and support.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL: Not applicable

FUNDING: This study received no specific funding from public, commercial, or not-for-profit funding agencies.

AI TOOL DECLARATION

The authors declare that no AI and related tools are used to write the scientific content of this manuscript.

DATA AVAILABILITY

Data will be available on request

REFERENCES

[1] Aaliyu AB, Musa AM, Oshanimi JA, Ibrahim HA and Oyewale AO (2008), Phytochemical analyses and mineral elements composition of some medicinal plants of northern Nigeria. Journ. Pharm. Sci., 7(1): 119–125.

[2] Salma sultana Sk, Banurekha J, Kumar M. Screening of anti colon cancer activity by using hydro alcoholic extract of Borassus Fabilifers hoot on swiss albino mice. Asian journal of Pharmaceutical and Clinical Research. 2025:18(7): 129-134

[3] Hoffman JM, Brooks EM, Mawe GM. Gastrointestinal motility mechanisms. J Vis Exp. 2011;78(3):2435.

[4] Truhwald S, Holzer P, Metzler H. Gastrointestinal dysfunction in critical illness. Intensive Care Med. 2000;56(34):1-8.

[5] Holger P, Schicho R, Holzer-Petsche U, Lippe I. The gut as a neurological organ. Wien Klin Wochenschr. 2001;113(4):647–600.

[6] Langley JN. The autonomic nervous system. Cambridge: Heffer; 1921.

[7] Camilleri M. Gastrointestinal motility disorders in neurologic disease. J Clin Invest. 2018;128(7):2981-2989.

[8] Huizinga JD, Lammers WJEP. Gut peristalsis mechanisms. Am J Physiol Gastrointest Liver Physiol. 2009;296(1):G1-G8.

[9] Spencer NJ, Hu H. Mechanisms of gastrointestinal motility. Neurogastroenterol Motil. 2020; 32(5): e13854.

[10] Farre R, Tack J. Pathophysiology of motility disorders. Nat Rev Gastroenterol Hepatol. 2013;10(8):455-465.